

**Quality ID #50: Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older**

- National Quality Strategy Domain: Person and Caregiver-Centered Experience and Outcomes
- Meaningful Measure Area: Management of Chronic Conditions

**2020 COLLECTION TYPE:**

**MIPS CLINICAL QUALITY MEASURES (CQMS)**

**MEASURE TYPE:**

Process – High Priority

**DESCRIPTION:**

Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months

**INSTRUCTIONS:**

This measure is to be submitted a minimum of **once per performance period** for patients seen during the performance period. This measure is appropriate for use in the ambulatory setting only. It is anticipated that Merit-based Incentive Payment System (MIPS) eligible clinicians who provide services for patients with the diagnosis of urinary incontinence will submit this measure.

**Measure Submission Type:**

Measure data may be submitted by individual MIPS eligible clinicians, groups, or third party intermediaries. The listed denominator criteria are used to identify the intended patient population. The numerator options included in this specification are used to submit the quality actions as allowed by the measure. The quality-data codes listed do not need to be submitted by MIPS eligible clinicians, groups, or third party intermediaries that utilize this modality for submissions; however, these codes may be submitted for those third party intermediaries that utilize Medicare Part B claims data. For more information regarding Application Programming Interface (API), please refer to the Quality Payment Program (QPP) website.

**DENOMINATOR:**

All female patients aged 65 years and older with a diagnosis of urinary incontinence

**Denominator Criteria (Eligible Cases):**

All female patients aged  $\geq 65$  years on date of encounter

**AND**

**Diagnosis for urinary incontinence (ICD-10-CM):** F98.0, N39.3, N39.41, N39.42, N39.43, N39.44, N39.45, N39.46, N39.490, N39.491, N39.492, N39.498, R32

**AND**

**Patient encounter during the performance period (CPT or HCPCS):** 99201, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99324, 99325, 99326, 99327, 99328, 99334, 99335, 99336, 99337, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, G0402

**AND NOT**

**DENOMINATOR EXCLUSION:**

**Hospice services utilized by patient any time during the measurement period:** G9694

**NUMERATOR:**

Patients with a documented plan of care for urinary incontinence at least once within 12 months

**Definition:**

**Plan of Care** – May include behavioral interventions (e.g., bladder training, pelvic floor muscle training, prompted voiding), referral to specialist, surgical treatment, reassess at follow-up visit, lifestyle interventions, addressing co-morbid factors, modification or discontinuation of medications contributing to urinary incontinence, or pharmacologic therapy.

**Numerator Options:**

***Performance Met:***

Urinary incontinence plan of care documented  
**(0509F)**

**OR**

***Performance Not Met:***

Urinary incontinence plan of care not documented,  
reason not otherwise specified **(0509F with 8P)**

**RATIONALE:**

A treatment option should be documented for the patient with incontinence.

**CLINICAL RECOMMENDATION STATEMENTS:**

All conservative management options used in younger adults can be used in selected frail, older, motivated people. This includes:

- Bladder retraining
- Pelvic muscle exercises including biofeedback and/or electro-stimulation (ICI) (Grade B)

Pharmacologic agents, especially oxybutynin and tolterodine, may have a small beneficial effect on improving symptoms of detrusor over activity in women. (ACOG) (Level A)

Oxybutynin and potentially other bladder relaxants can improve the effectiveness of behavioral therapies in frail older persons. (ICI) (Grade B)

**COPYRIGHT:**

Physician Performance Measures (Measures) and related data specifications have been developed by the PCPI® Foundation (PCPI®) and the National Committee for Quality Assurance (NCQA). These Measures are not a clinical guideline and does not establish a standard of medical care, and has not been tested for all potential applications. The Measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, eg, use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the Measures require a license agreement between the user and the PCPI® or NCQA. Neither the American Medical Association (AMA), nor the former AMA-convened Physician Consortium for Performance Improvement(R), PCPI, NCQA nor its members shall be responsible for any use of the Measure. © 2019 NCQA and PCPI® Foundation. All Rights Reserved.


THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND.

Limited proprietary coding is contained in the Measure specifications for user convenience. Users of proprietary code sets should obtain all necessary licenses from the owners of the code sets. NCQA disclaims all liability for use or accuracy of any CPT or other codes contained in the specifications.

CPT® contained in the Measure specifications is copyright 2004-2019 American Medical Association. LOINC® copyright 2004-2019 Regenstrief Institute, Inc. This material contains SNOMED Clinical Terms® (SNOMED CT®) copyright 2004-2019 International Health Terminology Standards Development Organisation. ICD-10 is copyright 2019 World Health Organization. All Rights Reserved.

**2020 Clinical Quality Measure Flow for Quality ID #50:  
Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older**

*Disclaimer: Refer to the measure specification for specific coding and instructions to submit this measure*



**SAMPLE CALCULATIONS:**

**Data Completeness=**  

$$\frac{\text{Performance Met (a=40 patients)} + \text{Performance Not Met (c=30 patients)}}{\text{Eligible Population / Denominator (d=80 patients)}} = \frac{70 \text{ patients}}{80 \text{ patients}} = 87.50\%$$

**Performance Rate=**  

$$\frac{\text{Performance Met (a=40 patients)}}{\text{Data Completeness Numerator (70 patients)}} = \frac{40 \text{ patients}}{70 \text{ patients}} = 57.14\%$$

\*See the posted measure specification for specific coding and instructions to submit this measure.

NOTE: Submission Frequency: Patient-Process

CPT only copyright 2019 American Medical Association. All rights reserved.  
The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification. v4

**2020 Clinical Quality Measure Flow Narrative for Quality ID #50:**  
**Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older**

**Disclaimer:** Refer to the measure specification for specific coding and instructions to submit this measure.

1. Start with Denominator
2. Check Female Patient Age:
  - a. If Female Patient Age is greater than or equal to 65 Years on Date of Encounter equals No during the measurement period, do not include in Eligible Population. Stop Processing.
  - b. If Female Patient Age is greater than or equal to 65 Years on Date of Encounter equals Yes during the measurement period, proceed to check Patient Diagnosis.
3. Check Patient Diagnosis:
  - a. If Diagnosis of Urinary Incontinence as Listed in the Denominator equals No, do not include in Eligible Population. Stop Processing.
  - b. If Diagnosis of Urinary Incontinence as Listed in the Denominator equals Yes, proceed to check Encounter Performed.
4. Check Encounter Performed:
  - a. If Encounter as Listed in the Denominator equals No, do not include in Eligible Population. Stop Processing.
  - b. If Encounter as Listed in the Denominator equals Yes, proceed to check Hospice Services Utilized Any Time During the Measurement Period
5. Check Hospice Services Utilized Any Time During the Measurement Period:
  - a. If Hospice Services Utilized Any Time During the Measurement Period equals No, include in Eligible Population.
  - b. If Hospice Services Utilized Any Time During the Measurement Period equals Yes, do not include in Eligible Population. Stop Processing.
6. Denominator Population
  - a. Denominator Population is all Eligible Patients in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d equals 80 patients in the Sample Calculation.
7. Start Numerator
8. Check Plan of Care for Urinary Incontinence Documented:
  - a. If Plan of Care for Urinary Incontinence Documented equals Yes, include in Data Completeness Met and Performance Met.

- b. Data Completeness Met and Performance Met letter is represented as Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a equals 40 patients in the Sample Calculation.
  - c. If Plan of Care for Urinary Incontinence Documented equals No, proceed to check Plan of Care for Urinary Incontinence Not Documented, Reason Not Otherwise Specified.
9. Check Plan of Care for Urinary Incontinence Not Documented Reason Not Otherwise Specified:
- a. If Plan of Care for Urinary Incontinence Not Documented, Reason Not Otherwise Specified equals Yes, include in Data Completeness Met and Performance Not Met.
  - b. Data Completeness Met and Performance Not Met letter is represented as Data Completeness in the Sample Calculation listed at the end of this document. Letter c equals 30 patients in the Sample Calculation.
  - c. If Plan of Care for Urinary Incontinence Not Documented, Reason Not Otherwise Specified equals No, proceed to check Data Completeness Not Met.
10. Check Data Completeness Not Met:
- a. If Data Completeness Not Met, the Quality Data Code or equivalent was not submitted. 10 patients have been subtracted from the Data Completeness Numerator in the Sample Calculation.

**SAMPLE CALCULATIONS:**

**Data Completeness=**

$$\frac{\text{Performance Met (a=40 patients)} + \text{Performance Not Met (c=30 patients)}}{\text{Eligible Population / Denominator (d=80 patients)}} = \frac{70 \text{ patients}}{80 \text{ patients}} = 87.50\%$$

**Performance Rate=**

$$\frac{\text{Performance Met (a=40 patients)}}{\text{Data Completeness Numerator (70 patients)}} = \frac{40 \text{ patients}}{70 \text{ patients}} = 57.14\%$$